



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N	A2	(11) International Publication Number: WO 99/63813 (43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/EP99/03883 (22) International Filing Date: 4 June 1999 (04.06.99) (30) Priority Data: 9812331.8 8 June 1998 (08.06.98) GB 9903669.1 17 February 1999 (17.02.99) GB (71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH). (71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT). (72) Inventors; and (75) Inventors/Applicants (for US only): ZURFLÜH, René [CH/CH]; Dachslenbergstrasse 54, CH-8180 Bülach (CH). LEADBITTER, Neil [GB/CH]; Therwilerstrasse 117, CH-4104 Oberwil (CH). (74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: FUNGICIDAL COMBINATIONS COMPRISING GLYOXALIC ACID METHYL ESTER-O-METHYLOXIME DERIVATIVES (57) Abstract <p>A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease an effective amount of a combination of a) 2-[α-{[(α-methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I) in association with b) a broad variety of other plant fungicides is particularly effective in combating or preventing diseases of crop plants. These combinations exhibit synergistic fungicidal activity.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

FUNGICIDAL COMBINATIONS COMPRISING GLYOXALIC ACID METHYL ESTER-O-METHYLOXIME DERIVATIVES


The present invention relates to novel fungicidal compositions for the treatment of phytopathogenic diseases of crop plants, especially phytopathogenic fungi, and to a method of combating phytopathogenic diseases on crop plants.

It is known that certain strobilurin derivatives have biological activity against phytopathogenic fungi, e.g. from EP-A-460575 where their properties and methods of preparation are described. On the other hand anilide, carbamate and aminoacid amide fungicides are widely known as plant fungicides for application in various crops of cultivated plants. However, crop tolerance and activity against phytopathogenic plant fungi do not always satisfy the needs of agricultural practice in many incidents and aspects.

It has now been found that the use of


- a) 2-[α -{[(α -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime, compound I (EP-460575)
in association with

- b) either a compound of formula IIA




an anilide of formula IIB (EP-545099)

- 2 -



wherein R₁ is fluorine or chlorine; or

a carbamate of formula IIC (WO-96/01256 and WO-96/01258)



wherein X is N or CH, and R₂ is 4-CH₃, 4-Cl or 2,4-dichloro; or


a compound IID (EP-278595)

methyl(2)-2-{2-[6-(trifluoromethyl)pyrid-2-yloxymethyl]-phenyl}-3-methoxyacrylate; or

a compound IIE (EP-477631)

(E)-N-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide; or


a compound of formula IIF (WO-95/21154)



wherein R₃ is methyl or ethyl; or

a (S)-valinamide of formula IIG (EP-398072, EP-610764, DE-4321897, WO-96/07638)

- 3 -




wherein R₄ is isopropyl, sec.-butyl or tert.-butyl, and

R₅ is 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl or β-naphthyl, and

wherein the asymmetric center is preferably (R); or

a (S)-valinamide of formula IIH (WO-94/25432, WO-96/04252)




wherein R₆ is isopropyl, sec.-butyl or tert.-butyl, R₇ is halogen, methyl or methoxy and n is 0,1 or 2; or

a compound IIJ (EP-596254)

N-methyl-2-[2-{α-methyl-3-(trifluoromethyl)benzyloximinomethyl}phenyl]-2-methoximinoacetamide; or

a compound of formula IIK (EP-381330)



wherein R₈ is halogen or C₁-C₄-alkyl, preferably chlorine; or


a compound II L

N-(3'-(1'-chloro-3-methyl 2'-oxopentane))-3,5-dichloro-4-methylbenzamide (EP-600629); or


a compound IIM (EP-551048 and WO 96/03044)

(S)-1-anilino-4-methyl-2-methylthio-4-phenylimidazolin-5-one; or

a compound of formula IIN (WO 98/25465)



a compound of formula IIP (WO98/20003)



a compound IIQ

N-methyl-2-[α -{[(α -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acidamide-O-methyloxime (EP 569384)

is particularly effective in combating or preventing fungal diseases of crop plants. These combinations exhibit synergistic fungicidal activity.

The combinations according to the invention may also comprise more than one of the active components b) , if broadening of the spectrum of disease control is desired.

The active ingredient combinations are effective against phytopathogenic fungi belonging to the following classes: Ascomycetes (e.g. *Venturia*, *Podosphaera*, *Erysiphe*, *Monilinia*, *Mycosphaerella*, *Uncinula*); Basidiomycetes (e.g. the genus *Hemileia*, *Rhizoctonia*, *Puccinia*); Fungi imperfecti (e.g. *Botrytis*, *Helminthosporium*, *Rhynchosporium*, *Fusarium*, *Septoria*, *Cercospora*, *Alternaria*, *Pyricularia* and *Pseudocercospora herpotrichoides*

(*Tapesia spp.*); Oomycetes (e.g. *Phytophthora*, *Peronospora*, *Bremia*, *Pythium*, *Plasmopara*).

Target crops for the areas of indication disclosed herein comprise within the scope of this invention e.g. the following species of plants: cereals (wheat, barley, rye, oats, rice, sorghum and related crops); beet (sugar beet and fodder beet); pomes, stone fruit and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (marrows, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocados, cinnamon, camphor); or plants such as maize, tobacco, nuts, coffee, sugar cane, tea, vines, hops, bananas and natural rubber plants, as well as ornamentals (flowers, shrubs, broad-leaved trees and evergreens, such as conifers). This list does not represent any limitation.

The combinations according to the present invention are particularly effective against *Phytophthora*, *Peronospora*, *Bremia*, *Pythium* and *Plasmopara*, in particular against pathogens of monocotyledonous plants such as cereals, including wheat and barley.

The amount of combination of the invention to be applied, will depend on various factors such as the compound employed, the subject of the treatment (plant, soil, seed), the type of treatment (e.g. spraying, dusting, seed dressing), the purpose of the treatment (prophylactic or therapeutic), the type of fungi to be treated and the application time.

Particularly preferred mixing partners of the compound I are those which comprise as component b) a compound IIB, IIG, IIH, IIK or IIL.

Another preferred mixing partners of the compound I are those which comprise as component b) a compound IIM, IIN, IIP or IIQ.

Another embodiment of the present invention is represented by those combination which comprise as component a) the compound I and as component b) a compound IIC, IID, IIE, IIF or IIJ.

Another combination is represented by the mixture comprising as component a) the compound I and as component b) the compound of the formula IIA.

It has been found that the use of compound I in combination with the compounds of formula II surprisingly and substantially enhances the effectiveness of the latter against fungi, and vice versa. Additionally, the method of the invention is effective against a wider spectrum of such fungi that can be combated with the active ingredients of this method when used solely.

The weight ratio of a):b) is so selected as to give a synergistic fungicidal action. In general the weight ratio of a) : b) is between 10 : 1 and 1 : 20. The synergistic action of the composition is apparent from the fact that the fungicidal action of the composition of a) + b) is greater than the sum of the fungicidal actions of a) and b).

Where the component b) is the compound IIA the weight ratio of a):b) is for example between 6:1 and 1:6, especially 2:1 and 1:2.

Where the component b) is a compound of formula IIB the weight ratio of a):b) is for example between 5:1 and 1:20, especially 2:1 and 1:20, and more preferably 1:1 to 1:10.

Where component b) is a compound of formula IIC, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IID, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IIE, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIF, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIG, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIH, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is the compound IIJ, the weight ratio of a) : b) is for example between 5:1 and 1:5, especially 3:1 and 1:3, and more preferably 2:1 and 1:2.

Where component b) is a compound of formula IIK, the weight ratio of a) : b) is for example between 5:1 and 1:20, especially 3:1 and 1:10, and preferably 2:1 and 1:5.

Where component b) is the compound IIL, the weight ratio of a) : b) is for example between 5:1 and 1:5, specially 2:1 and 1:2, and more preferably 1.5:1 and 1:1.5.

Where component b) is the compound IIM, the weight ratio of a) : b) is for example between 5:1 and 1:5, specially 2:1 and 1:2.

Where component b) is the compound IIN, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

Where component b) is the compound IIP, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

Where component b) is the compound IIQ, the weight ratio of a) : b) is for example between 6:1 and 1:6, specially 2:1 and 1:2.

The method of the invention comprises applying to the treated plants or the locus thereof in admixture or separately, a fungicidally effective aggregate amount of compound I and a compound of component b).

The term locus as used herein is intended to embrace the fields on which the treated crop plants are growing, or where the seeds of cultivated plants are sown, or the place where the seed will be placed into the soil. The term seed is intended to embrace plant propagating material such as cuttings, seedlings, seeds, germinated or soaked seeds.

The novel combinations are extremely effective on a broad spectrum of phytopathogenic fungi, in particular from the Fungi imperfecti and Oomycetes classes. Some of them have a systemic action and can be used as foliar and soil fungicides.

The fungicidal combinations are of particular interest for controlling a large number of fungi in various crops or their seeds, especially wheat, rye, barley, oats, rice, maize, lawns, cotton, soybeans, coffee, sugarcane, fruit and ornamentals in horticulture and viticulture, and in vegetables such as cucumbers, beans and cucurbits.

The combinations are applied by treating the fungi or the seeds, plants or materials threatened by fungus attack, or the soil with a fungicidally effective amount of the active ingredients.

The agents may be applied before or after infection of the materials, plants or seeds by the fungi.

The novel combinations are particularly useful for controlling the following plant diseases:

Erysiphe graminis in cereals,

Erysiphe cichoracearum and *Sphaerotheca fuliginea* in cucurbits,

Podosphaera leucotricha in apples,

Uncinula necator in vines,

Puccinia species in cereals,

Rhizoctonia species in cotton, rice and lawns,

Ustilago species in cereals and sugarcane,

Venturia inaequalis (scab) in apples,

Helminthosporium species in cereals,

Septoria nodorum in wheat,

Septoria tritici in wheat wheat,

Rhynchosporium secalis on barley

Botrytis cinerea (gray mold) in strawberries, tomatoes and grapes,

Cercospora arachidicola in groundnuts,

Peronospora tabacina on tobacco,

Bremia lactucae on lettuce,

Pythium debaryanum on sugar beet,

Pseudocercospora herpotrichoides (*Tapesia* spp.) in wheat and barley,

Pyrenophora teres in barley

Pyricularia oryzae in rice,

Phytophthora infestans in potatoes and tomatoes,

Fusarium and *Verticillium* species in various plants,

Plasmopara viticola in grapes,
Alternaria species in fruit and vegetables.

When applied to the plants the compound I is applied at a rate of 50 to 200 g/ha, particularly 75 to 150 g/ha, e.g. 75, 100, or 125g/ha, in association with 50 to 1500 g/ha, particularly 60 to 1000 g/ha, e.g. 75 g/ha, 80 g/ha, 100 g/ha, 125 g/ha, 150 g/ha, 175 g/ha, 200 g/ha, 300 g/ha, 500 g/ha, or 1000 g/ha of a compound of component b), depending on the class of chemical employed as component b). Where the component b) is the compound IIA for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIB for example 50 to 1500 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIC for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IID for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIE for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIF for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIG for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIH for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIJ for example 50 to 300 g a.i./ha is applied in association with the compound I. Where the component b) is a compound of formula IIK for example 20 to 2000 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIL for example 50 to 200 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIM for example 50 to 200 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIN for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIP for example 50 to 400 g a.i./ha is applied in association with the compound I. Where the component b) is the compound IIQ for example 50 to 400 g a.i./ha is applied in association with the compound I.

In agricultural practice the application rates depend on the type of effect desired, and range from 0.02 to 3 kg of active ingredient per hectare.

When the active ingredients are used for treating seed, rates of 0.001 to 50, and preferably from 0.01 to 10g per kg of seed are generally sufficient.

The invention also provides fungicidal compositions comprising the compound I and a compound of component b).

The composition of the invention may be employed in any conventional form, for example in the form of a twin pack, an instant granulate, a flowable or a wettable powder in combination with agriculturally acceptable adjuvants. Such compositions may be produced in conventional manner, e.g. by mixing the active ingredients with appropriate adjuvants (diluent or solvents and optionally other formulating ingredients such as surfactants). Suitable carriers and adjuvants may be solid or liquid and correspond to the substances ordinarily employed in formulation technology, such as, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binding agents or fertilizers. Such carriers are for example described in WO 96/22690.

Particularly formulations to be applied in spraying forms such as water dispersible concentrates or wettable powders may contain surfactants such as wetting and dispersing agents, e.g. the condensation product of formaldehyde with naphthalene sulphonate, an alkylarylsulphonate, a lignin sulphonate, a fatty alkyl sulphate, and ethoxylated alkylphenol and an ethoxylated fatty alcohol.

A seed dressing formulation is applied in a manner known per se to the seeds employing the combination of the invention and a diluent in suitable seed dressing formulation form, e.g. as an aqueous suspension or in a dry powder form having good adherence to the seeds. Such seed dressing formulations are known in the art. Seed dressing formulations may contain the single active ingredients or the combination of active ingredients in encapsulated form, e.g. as slow release capsules or microcapsules.

In general, the formulations include from 0.01 to 90% by weight of active agent, from 0 to 20% agriculturally acceptable surfactant and 10 to 99.99% solid or liquid adjuvant(s), the active agent consisting of at least the compound of formula I together with a compound of component b), and optionally other active agents, particularly guazatin and fenpiclonil. Concentrate forms of compositions generally contain in between about 2 and 80%, preferably between about 5 and 70% by weight of active agent. Application forms of formulation may for example contain from 0.01 to 20% by weight, preferably from 0.01 to 5% by weight of active agent.

Examples for specific formulations-combination are as disclosed e.g. in WO 96/22690, e.g. for wettable powders, emulsifiable concentrate, dusts, extruder granules, coated granules, suspension concentrate.

Slow Release Capsule Suspension

28 parts of a combination of the compound I and a compound of component b), or of each of these compounds separately, are mixed with 2 parts of an aromatic solvent and 7 parts of toluene diisocyanate/polymethylene-polyphenylisocyanate-mixture (8:1). This mixture is emulsified in a mixture of 1.2 parts of polyvinylalcohol, 0.05 parts of a defoamer and 51.6 parts of water until the desired particle size is achieved. To this emulsion a mixture of 2.8 parts 1,6-diaminohexane in 5.3 parts of water is added. The mixture is agitated until the polymerization reaction is completed.

The obtained capsule suspension is stabilized by adding 0.25 parts of a thickener and 3 parts of a dispersing agent. The capsule suspension formulation contains 28% of the active ingredients. The medium capsule diameter is 8-15 microns.

The resulting formulation is applied to seeds as an aqueous suspension in an apparatus suitable for that purpose.

Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

Biological Examples

A synergistic effect exists whenever the action of an active ingredient combination is greater than the sum of the actions of the individual components.

The action to be expected E for a given active ingredient combination obeys the so-called COLBY formula and can be calculated as follows (COLBY, S.R. "Calculating synergistic and antagonistic responses of herbicide combination". Weeds, Vol. 15, pages 20-22; 1967):

ppm = milligrams of active ingredient (= a.i.) per litre of spray mixture

X = % action by active ingredient I using p ppm of active ingredient

Y = % action by active ingredient II using q ppm of active ingredient.

According to Colby, the expected (additive) action of active ingredients I+II using p+q ppm

of active ingredient is $E = X + Y - \frac{X \cdot Y}{100}$

If the action actually observed (O) is greater than the expected action (E), then the action of the combination is superadditive, i.e. there is a synergistic effect.

Alternatively the synergistic action may also be determined from the dose response curves according to the so-called WADLEY method. With this method the efficacy of the a.i. is determined by comparing the degree of fungal attack on treated plants with that on untreated, similarly inoculated and incubated check plants. Each a.i. is tested at 4 to 5 concentrations. The dose response curves are used to establish the EC90 (i.e. concentration of a.i. providing 90% disease control) of the single compounds as well as of the combinations (EC 90_{observed}). The thus experimentally found values of the mixtures at a given weight ratio are compared with the values that would have been found were only a complementary efficacy of the components was present (EC 90 (A+B)_{expected}). The EC90 (A+B)_{expected} is calculated according to Wadley (Levi et al., EPPO- Bulletin 16, 1986, 651-657):

$$\text{EC 90 (A+B)}_{\text{expected}} = \frac{a + b}{\frac{a}{\text{EC90 (A)}_{\text{observed}}} + \frac{b}{\text{EC90 (B)}_{\text{observed}}}}$$

wherein a and b are the weight ratios of the compounds A and B in the mixture and the indexes (A), (B), (A+B) refer to the observed EC 90 values of the compounds A, B or the given combination A+B thereof. The ratio EC90 (A+B)_{expected} / EC90 (A+B)_{observed} expresses the factor of interaction (F). In case of synergism, F is >1.

Example B-1: Residual-protective action against *Venturia inaequalis* on apples

Apple cuttings with 10-20 cm long fresh shoots are sprayed to drip point with an aqueous spray mixture prepared from a wettable powder formulation of the active ingredient mixture and infected 24 hours later with a conidia suspension of the fungus. The plants are incubated for 5 days at 90-100 % relative humidity and stood in a greenhouse for a further 10 days at 20-24°C. Fungus infestation is evaluated 12 days after infection.

Example B-2(a): Action against *Botrytis cinerea* on apple fruits

Artificially damaged apples are treated by dropping a spray mixture of the active ingredient mixture onto the damage sites. The treated fruits are then inoculated with a spore suspension of the fungus and incubated for one week at high humidity and about 20°C. The fungicidal action of the test compound is derived from the number of damage sites that have begun to rot.

Example B-2(b): Action against *Botritis cinerea* on tomatoes

4 week old tomato plants cv. "Roter Gnom" were treated with the formulated testcompound in a spray chamber. Two days after application the tomato plants were inoculated by spraying a spore suspension on the test plants. After an incubation period of 4 days at 20°C and 95% relative humidity in a growth chamber the disease incidence was assessed.

Example B-2(c): Action against *Botritis cinerea* on grapes

5 week old grape seedlings cv. "Gutedel" were treated with the formulated testcompound in a spray chamber. Two days after application the grape plants were inoculated by spraying a spore suspension on the test plants. After an incubation period of 4 days at 21°C and 95% relative humidity in a greenhouse the disease incidence was assessed.

Example B-3: Action against *Podosphaera leucotricha* on apple shoots

Apple cuttings with about 15 cm long fresh shoots are sprayed with a spray mixture of the active ingredient mixture. The treated plants are infected 24 hours later with a conidia suspension of the fungus and placed in a climatic chamber at 70 % relative humidity and 20°C. Fungus infestation is evaluated 12 days after infection.

Example B-4: Action against *Drechslera teres* on barley

10-day-old barley plants of the "Golden Promise" variety are sprayed with a spray mixture of the active ingredient mixture. The treated plants are infected 24 hours later with a conidia suspension of the fungus and incubated in a climatic chamber at 70 % relative humidity and 20-22°C. Fungus infestation is evaluated 5 days after infection.

Example B-5 : Efficacy against *Erysiphe graminis* f.sp. *tritici* on wheat

Five to ten wheat seeds c.v. "Arina" are sown in plastic pots of 7 cm diameter and grown for 7 to 12 days at 20°C, 50-70% rH. When the primary leaves have fully expanded, the plants

are spray treated with aqueous spray liquors containing the single compounds, or mixtures thereof (hereinafter a.i.). All compounds are used as experimental or commercially available formulations, combinations are applied as tank mixtures. The application comprises foliar spraying to near runoff (three pots per treatment). 24 hours after the application or 24 hours before application, the plants are inoculated in a settling tower with fresh spores of *Erysiphe graminis* f. sp. *tritici*. The plants are then incubated in a growth chamber at 20°C, 60% rH. Seven days after the inoculation, the percentage of infection on primary leaves is evaluated. The efficacy of the a.i. is determined by comparing the degree of fungal attack on treated plants with that on untreated, similarly inoculated and incubated check plants. Each a.i. is tested at 4 to 5 concentrations. The synergy factor is calculated according to the COLBY method.

Example B-7: Activity against Uncinula necator

Grape plants, 4 weeks old (4-5 leaves), are sprayed to near run off with a suspension containing 250 mg/l of active ingredient. The deposit is then allowed to dry. One day later, the treated plants are inoculated by dusting freshly harvested conidia over the test plants; then the plants were incubated in a growth chamber for 10-14 days at +22°C and 70% r.h. The efficacy of the test compounds is determined by comparing the degree of fungal attack with that on untreated, similarly inoculated check plants.

The mixtures according to the invention exhibit good activity in these Examples.

Example B-8: Activity against Plasmopara viticola in grapevines

Grapevine seedlings at the 4- to 5-leaf stage are sprayed to drip point with an aqueous spray mixture prepared with a wettable powder of the active ingredient mixture (0.02% of active ingredient) and, 24 hours later, infected with a sporangia suspension of the fungus. The fungus infestation is assessed 6 days after infection, during which time a relative atmospheric humidity of 95 to 100% and a temperature of 20°C are maintained.

Example B-9: Activity against Phytophthora infestans in tomatoes

a) Curative action

Tomato plants cv. "Roter Gnom" are grown for three weeks and then sprayed with a zoospore suspension of the fungus and incubated in a cabin at 18 to 20°C and saturated atmospheric humidity. The humidification is interrupted after 24 hours. After the plants have dried, they are sprayed with a mixture which comprises the active ingredient formulated as a

wettable powder at a concentration of 200ppm. After the spray coating has dried, the plants are returned to the humid chamber for 4 days. Number and size of the typical foliar lesions which have appeared after this time are used as a scale for assessing the efficacy of the test substances.

b) Preventive-systemic action

The active ingredient which is formulated as a wettable powder is introduced, at a concentration of 60ppm (relative to the soil volume), onto the soil surface of three-week-old tomato plants cv. "Roter Gnom" in pots. After an interval of three days, the underside of the leaves is sprayed with a zoospore suspension of *Phytophthora infestans*. They are then kept for 5 days in a spray cabin at 18 to 20°C and saturated atmospheric humidity. After this time, typical foliar lesions appear whose number and size are used for assessing the efficacy of the test substances.

Example B-10: Activity against *Phytophthora* in potato plants

a) Residual-protective action

2-3 week old potato plants (Bintje variety) are grown for 3 weeks and then sprayed with a spray mixture (0.02% of active ingredient) prepared with a wettable powder of the active ingredient. After 24 hours, the treated plants are infected with a sporangia suspension of the fungus. The fungus infestation is assessed after the infected plants have been incubated for 5 days at a relative atmospheric humidity of 90-100% and 20°C.

b) Systemic action

A spray mixture (0.002% of active ingredient based on the soil volume) prepared with a wettable powder of the active ingredient is poured next to 2-3 week old potato plants (Bintje variety) which have been grown for 3 weeks. Care is taken that the spray mixture does not come into contact with the aerial parts of the plants. After 48 hours, the treated plants are infected with a sporangia suspension of the fungus. Fungus infestation is assessed after the infected plants have been incubated for 5 days at a relative atmospheric humidity of 90-100% and 20°C.

The efficacy of the test combinations and the single active ingredients in the above tests is determined by comparing the degree of fungal attack with that on untreated, similarly inoculated check plants.

The mixtures according to the invention exhibit good activity in these Examples.


WHAT IS CLAIMED IS:

1. A method of combating phytopathogenic diseases on crop plants which comprises applying to the crop plants or the locus thereof being infested with said phytopathogenic disease, in admixture or separately, an effective amount of a combination of a first component


a) 2-[α -{[(α -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acid methyl ester-O-methyloxime (I)

in association with

b) a second component selected from either
a compound of formula IIA




an anilide of formula IIB



wherein R₁ is fluorine or chlorine; or

a carbamate of formula IIC



wherein X is N or CH, and R₂ is 4-CH₃, 4-Cl or 2,4-dichloro; or


a compound IID

methyl(2)-2-[2-[6-(trifluoromethyl)pyrid-2-yloxymethyl]-phenyl]-3-methoxyacrylate; or

a compound IIE


(E)-N-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide; or

a compound of formula IIF



wherein R₃ is methyl or ethyl; or

a (S)-valinamide of formula IIG




wherein R₄ is isopropyl, sec.-butyl or tert.-butyl, and

R₅ is 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl or β-naphthyl, and

wherein the asymmetric center is preferably (R); or

a (S)-valinamide of formula IIH




wherein R₆ is isopropyl, sec.-butyl or tert.-butyl, R₇ is halogen, methyl or methoxy and

n is 0, 1 or 2; or

a compound IIJ

N-methyl-2-[2-(α -methyl-3-(trifluoromethyl)benzyloximinomethyl)phenyl]-2-methoximinoacetamide; or

a compound of formula IIK



wherein R_8 is halogen or C_1 - C_4 -alkyl, preferably chlorine; or


a compound III

N-(3'-(1'-chloro-3-methyl 2'-oxopentane))-3,5-dichloro-4-methylbenzamide; or


a compound IIM

(S)-1-anilino-4-methyl-2-methylthio-4-phenylimidazolin-5-one; or

a compound of formula IIN



a compound of formula IIP



a compound IIQ

N-methyl-2-[α -{[(α -methyl-3-trifluoromethyl-benzyl)imino]-oxy}-o-tolyl]-glyoxalic acidamide-O-methyloxime.

2. A method according to claim 1 wherein the components b) is selected from the compounds of formula IIB, IIG, IIH, IIK and IIL.
3. A method according to claim 1 wherein the component b) is selected from the group comprising IIC, IID, IIE, IIF and IIJ.
4. A method according to claim 1 wherein the component b) is a compound of formula IIA.
5. A method according to claim 1 wherein the component b) is selected from the group comprising IIM, IIN, IIP or IIQ.
6. A method according to any of claims 1 to 5 wherein the components a) and b) are applied in a quantity producing a synergistic disease controlling effect, specially a fungicidal effect.
7. A fungicidal composition comprising a fungicidally effective combination of
 - a) a compound I according to claim 1
in association with
 - b) either a compound IIA, or
a compound of formula IIB, or
a compound of formula IIC, or
a compound IID, or
a compound IIE, or
a compound of formula IIF, or
a compound of formula IIG, or
a compound of formula IIH, or
a compound IIJ, or
a compound of formula IIK, or
a compound IIL, or

a compound of formula IIM, or
a compound of formula IIN, or
a compound of formula IIP, or
a compound IIQ
as defined in claim 1.

8. A composition according to claim 7 the weight ratio of a) to b) is between 10 : 1 and 1 : 20.

9. A composition according to claim 7 wherein the component b) is selected from the group comprising IIB, IIG, IIH, IIK and IIL.

10. A composition according to claim 7 wherein the component b) is selected from the group comprising IIC, IID, IIE, IIF and IIJ.

11. A composition according to claim 7 wherein the component b) is a compound of formula IIA.

12. A composition according to claim 7 wherein the component b) is selected from the group comprising IIM, IIN, IIP and IIQ.